

United States Patent Application

of

Dr. Uwe Heinelt
Dr. Hans-Jochen Lang
Dr. Klaus Wirth
and
Dr. Hans-Willi Jansen

for

**Substituted heterocyclo-norbornylamino derivatives, processes for
their preparation, their use as medicaments or diagnostics, and
medicaments comprising them**


FILED

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[001] This application claims the benefit of the filing date of German Patent Application Number 10063294.7-44, filed on December 19, 2000, which application is hereby incorporated by reference.

[002] One embodiment of the invention relates to substituted heterocyclo-norbornylamino derivatives having: a) an exo-configuration nitrogen and an endo-fused five-membered or six-membered ring of the formula (I), or b) an exo-configuration nitrogen and an exo-fused five-membered or six-membered ring of the formula (I a),



wherein:

[003] A is (C₁-C₄)- alkylene;

[004] T is (C₁-C₄)- alkyl or H;

[005] B is a saturated or unsaturated carbon five-membered or six-membered ring, which is unsubstituted or is substituted, having 1-3 substituents chosen from oxo, hydroxyl, (C₁-C₄)- alkoxy, and (C₁-C₄)- alkyl;

[006] Het is a 5- or 6-membered, saturated or unsaturated, heterocycle that contains up to four identical or different heteroatoms chosen from O, S, N, and Se;

[007] R is OH, F, Cl, Br, I, CN, NO₂, phenyl, CO₂R₁, (C₁-C₄)- alkyl, (C₁-C₄)- alkoxy, amino, (C₁-C₄)- alkylamino, di(C₁-C₄)- alkylamino, or amino-(C₁-C₄)- alkyl,

[008] wherein the alkyl radicals are unsubstituted or are completely or partly substituted by fluorine;

[009] R₁ is H or (C₁-C₄)-alkyl, which is unsubstituted or completely or partly substituted by fluorine;

[010] n0, 1, 2, 3 or 4,

[011] wherein, if n = 2, 3 or 4, the substituents R are chosen independently of one another;

[012] and their pharmaceutically tolerable salts or trifluoroacetates.

[013] Examples of compounds of the invention include those compounds having an exo-configuration nitrogen and endo-fused carbon five-membered or six-membered ring of the formula (I) and those compounds having an exo-configuration nitrogen and exo-fused carbon five-membered or six-membered ring of the formula (I a), wherein:

[014] A is (C₁-C₂)- alkylene;

[015] T is H or methyl;

[016] B is a saturated or unsaturated carbon five-membered or six-membered ring;

[017] Het is a 5- or 6-membered, saturated or unsaturated, heterocycle that contains up to three identical or different heteroatoms chosen from O, S, and N;

[018] R is F, Cl, Br, iodine, amino, hydroxymethyl, OH, phenyl, CO₂R₁, (C₁-C₄)- alkyl, or (C₁-C₄)- alkoxy,

[019] wherein the alkyl radicals are unsubstituted or completely or partly substituted by fluorine;

[020] R₁ is H or (C₁-C₄)-alkyl, wherein the alkyl radical is unsubstituted or completely or partly substituted by fluorine;

[021] n0, 1, 2 or 3,

[022] wherein, if n = 2 or 3, the corresponding substituents R are chosen independently of one another;

[023] and their pharmaceutically tolerable salts or trifluoroacetates.

[024] Further examples of compounds of the invention include those compounds having an exo-configuration nitrogen and endo-fused carbon five-membered or six-membered ring of the formula (I) and those compounds having an

exo-configuration nitrogen and exo-fused carbon five-membered or six-membered ring of the formula (I a), wherein:

[025] A is (C₁-C₂)- alkylene;

[026] T is hydrogen;

[027] B is a saturated or unsaturated carbon five-membered or six-membered ring;

[028] Het is a 5- or 6-membered, saturated or unsaturated, heterocycle that contains up to two identical or different heteroatoms chosen from O, S, and N ;

[029] R is F, Cl, Br, (C₁-C₄)- alkoxy or (C₁-C₄)- alkyl,

[030] wherein the alkyl radicals are unsubstituted or completely or partly substituted by fluorine;

[031] n is 0, 1 or 2, wherein, if n = 2, the corresponding substituents R are chosen independently of one another;

[032] and their pharmaceutically tolerable salts or trifluoroacetates.

[033] Even more examples of the compounds of the invention include those compounds having an exo-configuration nitrogen and endo-fused carbon five-membered or six-membered ring of the formula (I) and those compounds having an exo-configuration nitrogen and exo-fused carbon five-membered ring of the formula (I a) such as, for example:

exo/exo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
 (rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
 (+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
 (-)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine,
 (rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,
 (+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,
 (-)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyrazin-2-ylmethylamine,
 exo/endo-(decahydro-1,4-methanonaphthalen-2-yl)pyrazin-2-ylmethylamine,
 exo/endo-(octahydro-4,7-methanoinden-5-yl)thiophen-2-ylmethylamine,
 exo/endo-(octahydro-4,7-methanoinden-5-yl)thiophen-3-ylmethylamine,

[illegible]

LAW OFFICES
EGAN, HENDERSON,
LABOW, GARRETT,
DUNNER, L.L.P.
101 STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

- 4 -


and their pharmaceutically tolerable salts or trifluoroacetates.

[035] Suitable acid addition salts of the compounds of the invention include salts of pharmacologically tolerable acids, for example halides (such as hydrochlorides), lactates, sulfates, citrates, tartrates, acetates, phosphates, methylsulfonates, p-toluenesulfonates, adipates, fumarates, gluconates, glutamates, glycerophosphates, maleates, and pamoates. This group also exemplifies some of the physiologically acceptable anions, in addition to trifluoroacetates.

[036] If a compound of the formula (I) or (I a) contains one or more asymmetric centers, these centers can have either the S or the R configuration. The compounds can be present as optical isomers, diastereomers, racemates, or mixtures thereof. However, the amino substituent on the norbornyl system must be exo and the ring endo- or exo-fused



[037] The alkyl or alkylene radicals mentioned can be either straight-chain or branched.

[038] Suitable heterocycles include, *inter alia*:




[039] Another embodiment of the invention relates to a process for the preparation of the compounds of the formula (I) or (I a), which comprises

[040] a) reacting a compound of the formula (II) or (II a)





[041] with a compound of the formula (III)



[042] in the presence of suitable reductants and possibly also Lewis acids directly to give compounds of the formula (I) or (I a),

[043] wherein T, B, Het and R_n have the meanings indicated above; independently of one another A' corresponds to a bond or (C₁-C₃)- alkyl and A'' corresponds to H or (C₁-C₃)- alkyl; and A' and A'', together with the carbon atom of the carbonyl group, represent as many carbon atoms as A represents in formula (I) or (I a); or

[044] b) isolating an intermediate of the formula (IV) or (IV a), formed from reacting compounds of the formulae (II), or (II a), with a compound of formula (III), and then converting the intermediate of the formula (IV) or (IV a) into the compounds of the formula (I) or (I a) by using suitable reductants,




LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[045] wherein T# is a free electron pair or (C₁-C₄)-alkyl. When T# is (C₁-C₄)-alkyl, an iminium ion is formed, to which a counterion is assigned, such as, for example, chloride or tosylate,

or



[046] c) reacting a compound of the formula (II) or (II a) with an alkylating agent of the formula (V),



[047] wherein U is a nucleophilically substitutable group, such as for example a halogen, alkylsulfonates, or arylsulfonates, including Cl, Br, I, mesylate, or tosylate, and the other radicals are defined as described above. Here, the carbon atom to which U is bonded corresponds to the carbon atom of the carbonyl group of the compound of formula (III),

or

[048] d) reducing carboxamides of the formula (VI) or (VI a) to the corresponding amines,



[049] wherein A* corresponds to a bond or (C₁-C₃)-alkyl and the other radicals are defined as described above,

or


[050] e) alkylating compounds of the formula (I) or (I a) , in which T corresponds to hydrogen, using alkylating agents of the formula (VII),




[051] wherein T* is(C₁-C₄)- alkyl and U has the meaning described above, so that tertiary amines result from this reaction;

or

[052] f) reacting a dicyclopentadienylplatinum complex of the formula (VIII)



[053] with amines of the formula (IX),



[054] and subsequently reducing the intermediate formed to compounds of the formula (I) or (I a) (J. K. Stille and D. B. Fox, JACS 92:1274 (1970)),

[055] wherein T, R_n and Het have the meanings indicated above; independently of one another, A' corresponds to a bond or (C₁-C₃)-alkyl and A'' to H or (C₁-C₃)-alkyl; and A' and A'', together with the carbon atom to which the nitrogen atom is bonded, represent as many carbon atoms as A represents in formula (I).

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[056] The compounds of formula (I) or (I a) may be optionally converted into the pharmaceutically tolerable salt or trifluoroacetate.

[057] It has already been proposed that phenylalkyl-substituted norbornylamino derivatives are effective inhibitors of the sodium-proton exchanger, subtype 3 (NHE3). In this case, it appears that, of several stereoisomers, the compounds having an exo/endo-configuration octahydro-4,7-methanoinden-5-ylamine unit, wherein the nitrogen is exo and the five-membered ring is endo-fused, are active NHE3 inhibitors. Substances having an exo/exo-configuration octahydro-4,7-methanoinden-5-ylamine unit likewise showed marked NHE3-inhibiting action, while the corresponding endo/endo and endo/exo-derivatives were markedly less active on the NHE3 (German published application 199 60 204 A1 - HMR 99/L 073).

[058] Surprisingly, it has now been found that the aromatic moiety of the phenylalkyl substituents can be substituted by heteroaromatic rings producing NHE3-inhibiting activity.

[059] The relatively long-known inhibitors of the sodium/proton exchanger, subtype 3 disclosed in EP-A 825 178 (HOE 96/F226) represent relatively polar structures and correspond to the acylguanidine type of compounds (J.-R. Schwark et al. Eur. J. Physiol (1998) 436:797). In contrast, the compounds according to the invention are surprisingly lipophilic substances that are not of the acylguanidine type. See also the proposed compounds of the phenylalkyl norbornylamine type (DE 199 60 204.2 - HMR 99 / L 073). Squalamine and the above phenylalkyl norbornylamino derivatives are only the fourth substance class of NHE3 inhibitors known hitherto (M. Donowitz *et al.*, Am. J. Physiol. 276 (Cell Physiol. 45): C136-C144). Additionally, squalamine does not achieve its maximum potency immediately, but only after approximately one hour. Compared with the above acylguanidines, the compounds of the invention are distinguished by their superior ability to cross the membrane and, compared with squalamine, by a more rapid onset of action.

0
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2

[060] NHE3 is found in the body of various species, mostly in the bile, the intestine, and in the kidney (Larry Fliegel *et al.*, Biochem. Cell. Biol. 76: 735-741 (1998)), but can also be detected in the brain (E. Ma *et al.*, Neuroscience 79: 591-603).

[061] The compounds of the formula (I) or (I a) according to the invention are suitable for use as antihypertensives for the treatment of primary and secondary hypertension.

[062] Moreover, these compounds, on their own or in combination with NHE inhibitors of other subtype specificity, can protect acutely or chronically oxygen-deficiently-supplied organs by reducing or preventing ischemically induced damage. Thus, the compounds of the invention are suitable as pharmaceuticals, in the treatment of acute or chronic kidney failure and during surgical interventions such as in organ transplantation of the kidney and liver, where the compounds can be used both for the protection of the organs in the donor before and during removal and for the protection of removed organs, (for example during treatment with or storage in physiological bath fluids, and during transfer to the recipient's body). Ischemically induced damage to the intestine can also be avoided.

[063] Corresponding to the protective action against ischemically induced damage, the compounds of the invention are also potentially suitable as pharmaceuticals for the treatment of ischemia of the nervous system, including the CNS, where they are suitable, for example, for the treatment of stroke or cerebral odema. Moreover, the compounds of the formula (I) or (I a) according to the invention are likewise suitable for the treatment of forms of shock, such as, for example, allergenic, cardiogenic, hypovolemic, and bacterial shock.

[064] Furthermore, the compounds of the invention induce an improvement in the respiratory drive and are therefore used for the treatment of respiratory conditions in the following clinical conditions and diseases: disturbed central respiratory drive (e.g.

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

central sleep apnea, sudden infant death, post operative hypoxia), muscular-related respiratory disorders, respiratory disorders after long-term ventilation, respiratory disorders during adaptation in a high mountain area, obstructive and mixed forms of sleep apnea, acute and chronic lung diseases with hypoxia and hypocapnia.

[065] The compounds of the invention additionally increase the muscle tone of the upper airways, so that snoring is suppressed.

[066] A combination of an NHE inhibitor with a carboanhydrase inhibitor, (e. g. acetazolamide) where the latter produces metabolic acidosis and thereby even increases the respiratory activity, proves to be advantageous as a result of increased activity and decreased use of active compound.

[067] It has been shown that the compounds according to the invention have a mild laxative action and accordingly can be used advantageously as laxatives or, in the case of threatening intestinal blockage, for the prevention of ischemic damage that accompanies blockages in the intestinal region.

[068] Furthermore, the use of the compounds of the formula (I) or (I a) according to the invention makes it possible to prevent gallstone formation.

[069] The compounds of the formula (I) or (I a) according to the invention additionally show an action against ectoparasites.

[070] Moreover, the compounds of the formula (I) or (I a) according to the invention can exert an inhibitory action on the proliferation of cells, for example fibroblast cell proliferation and the proliferation of the smooth vascular muscle cells. The compounds of the formula (I) or (I a) are therefore suitable as valuable therapeutics for diseases in which cell proliferation is a primary or secondary cause, and can therefore be used as antiatherosclerotics, as agents against diabetic late complications, cancers,

202-408-4000

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

fibrotic disorders (such as, for example, pulmonary fibrosis, liver fibrosis, or kidney fibrosis), endothelial dysfunction, and organ hypotrophy and hyperplasia, such as, for example, prostate hyperplasia or prostate hypertrophy.

[071] The compounds according to the invention are effective inhibitors of the cellular sodium/proton antiporter, which is raised in numerous disorders (essential hypertension, atherosclerosis, diabetes etc.) even in those cells that are easily accessible to measurement, such as, for example, erythrocytes, platelets or leukocytes. The compounds according to the invention are therefore suitable as outstanding and simple scientific tools, for example in their use as diagnostics for the determination and differentiation of certain forms of hypertension, atherosclerosis, diabetes, proliferative disorders, etc. Moreover, the compounds of the formula (I) or (I a) are suitable for preventive therapy to prevent the development of high blood pressure, for example of essential hypertension.

[072] It has moreover been found that NHE inhibitors exhibit a favorable influence on the serum lipoproteins. It is generally recognized that for the formation of arteriosclerotic vascular changes, for example in coronary heart disease, excessively high blood lipid values, 'hyperlipoproteinemias', are an important risk factor. The lowering of increased serum lipoproteins is therefore of extreme importance for the prophylaxis and the regression of atherosclerotic changes. The compounds according to the invention can therefore be used for the prophylaxis and regression of atherosclerotic changes by excluding a causal risk factor. With this protection of the vessels against the syndrome of endothelial dysfunction, compounds of the formula (I) or (I a) are valuable pharmaceuticals for the prevention and treatment of coronary vascular spasms, atherogenesis and atherosclerosis, left-ventricular hypertrophy and dilated cardiomyopathy, and thrombotic disorders.

[073] The compounds mentioned are therefore advantageously used for the production of a medicament for the prevention and treatment of sleep apneas and

CONFIDENTIAL

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N.W.
WASHINGTON, DC 20005
202-408-4000

muscular-related respiratory disorders; for the production of a medicament for the prevention and treatment of snoring, for the production of a medicament for lowering the blood pressure, for the production of a medicament having laxative action for the prevention and treatment of intestinal blockages; for the production of a medicament for the prevention and treatment of disorders that are induced by ischemia and reperfusion of central and peripheral organs, such as acute kidney failure, stroke, endogenous states of shock, intestinal disorders, etc; for the production of a medicament for the treatment of hypercholesterolemia; for the production of a medicament for the prevention of atherogenesis and of athereosclerosis; for the production of a medicament for the prevention and treatment of diseases which are induced by raised cholesterol levels; for the production of a medicament for the prevention and treatment of diseases which are induced by endothelial dysfunction; for the production of a medicament for the treatment of attack by ectoparasites; for the production of a medicament for the treatment of the diseases mentioned in combination with blood pressure-lowering substances, for example angiotensin-converting enzyme (ACE) inhibitors and angiotensin receptor antagonists. A combination of an NHE inhibitor of the formula (I) or (I a) with an active compound lowering the blood lipid level, such as an HMG-CoA reductase inhibitor (e.g. Lovastatin or Pravastatin) proves to be a favorable combination having intensified action and decreased use of active substance. In this case, the HMG-CoA reductase inhibitor produces a hypolipodemic effect and thereby increases the hypolipodemic properties of the NHE inhibitor of the formula (I) or (I a).

[074] The administration of sodium/proton exchange inhibitors of the formula (I) or (I a) as novel pharmaceuticals for lowering raised blood lipid levels, and the combination of sodium/proton exchange inhibitors with pharmaceuticals having a blood pressure-lowering and/or hypolipodemic action is claimed.

[075] Pharmaceuticals containing a compound of formula (I) or (I a) can be administered orally, parenterally, intravenously, rectally, or by inhalation. The method of administration depends on the particular clinical picture of the disorder. The compounds

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

(I) or (I a) can be used on their own or together with pharmaceutical excipients, namely both in veterinary and in human medicine.

[076] The person skilled in the art is familiar on the basis of his/her expert knowledge with excipients that are suitable for the desired pharmaceutical formulation. In addition to solvents, gel formers, suppository bases, tablet excipients, and other active compound carriers, it is possible to use, for example, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents, preservatives, solubilizers, and/or colorants.

[077] For a form suitable for oral administration, the active compounds are ~~mixed with the additives suitable~~ therefor, including excipients, stabilizers, or inert diluents, and brought by means of the customary methods into the suitable administration forms, including tablets, coated tablets, hard gelatin capsules, aqueous, alcoholic, or oily solutions. Inert carriers that can be used include, for example, gum arabic, magnesia, magnesium carbonates, potassium phosphates, lactose, glucose, starch, or cornstarch. Preparation can be carried out either as dry or as moist granules. Possible oily excipients or solvents include, for example, vegetable or animal oils, such as, for example, sunflower oil or cod-liver oils.

[078] For subcutaneous or intravenous administration, the active compounds are brought into solution, suspension, or emulsion, if desired with the substances customary therefor such as, for example, solubilizers, emulsifiers, or further excipients. Suitable solvents include, for example, water, physiological saline solution, or alcohols (e.g. ethanol, propanol, glycerol), and sugar solutions such, for example, as glucose or mannitol solutions, or alternatively a mixture of the different solvents mentioned.

[079] Pharmaceutical formulations suitable for administration in the form of aerosols or sprays include, for example, solutions, suspensions, or emulsions of the active compound of the formula (I) or (I a) in a pharmaceutically innocuous solvent, such as, ethanol or water, or a mixture of such solvents.

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[080] If required, the formulation can also contain other pharmaceutical excipients such as, for example, surfactants, emulsifiers and stabilizers, and a propellant. Such a preparation may contain the active compound in a concentration of approximately 0.1 to 10 % by weight; and sometimes approximately 0.3 to 3 % by weight.

[081] The dose of the active compound of the formula (I) or (I a) to be administered and the frequency of administration depend on the potency and duration of action of the compounds used; on the nature and severity of the disease to be treated; and on the sex, age, weight, and individual responsiveness of the mammal to be treated.

[082] On average, the daily dose of a compound of the formula (I) or (I a) in the case of a patient weighing approximately 75 kg may be at least 0.001 mg/kg, sometimes 1 - 10 mg/kg, and at most 100 mg/kg, of bodyweight. In acute episodes of the diseases, even higher and more frequent doses may also be necessary, e.g. up to 4 individual doses per day, for instance in the case of i.v. administration, for example in the case of an infarct patient in the intensive care unit, up to 200 mg per day may be necessary.

Experimental section:

[083] Abbreviations used:

CI	chemical ionization
DIP	diisopropyl ether
EA	ethyl acetate
ES	electrospray
FAB	fast atom bombardment
h	hours

bp	boiling point
MgSO ₄	magnesium sulfate
m.p.	melting point
MS	mass spectrum
NaOH	sodium hydroxide solution
RP	reversed phase

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

HCl	hydrochloric acid
HPLC-RT	HPLC retention time

THF	tetrahydrofuran
TFA	trifluoroacetic acid

EXAMPLES:

[084] If not described otherwise, the examples mentioned here are racemates.

[085] The HPLC or LCMS conditions used for characterization were as follows: HPLC and HPLC-MSD systems from Agilent Technologies of the series 1100 with DAD detector, Merck Purospher column (3 μ , 2 x 55 mm), column temperature: 30°C, wavelength: 220 nm, flow: 0.5 ml/min, gradient: from 95% water (0.05% TFA)/5% acetonitrile in 4 min to 5% water (0.05% TFA)/95% acetonitrile, then the column was kept at 5% water (0.05% TFA)/95% acetonitrile for 1.5 min.


[086] The values marked by * were determined under the following conditions: HPLC-MSD system from Agilent Technologies of the series 1100 with DAD detector, Nucleosil column (C-18, 5 μ , 4x125 mm), column temperature: 40°C, wavelength: 220 nm, flow: 0.65 ml/min, gradient: 95% water [water/acetonitrile 9:1 with 0.1% TFA]/5% acetonitrile [water/acetonitrile 1:9 with 0.1% TFA] for 2 min, then to 5% water/95% acetonitrile in 10 min, then the column was kept at 5% water/95% acetonitrile for 5 min.

[087] For the characterization of the final compounds, the HPLC retention time and the result of the mass-spectroscopic investigation, which was carried out separately, are given.

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[088] Example 1: (rac)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine hydrochloride



[089] a) exo/endo-3a,4,5,6,7,7a-hexahydro-1H-4,7-methanoinden-5-ylamine and exo/endo-(3a,4,5,6,7,7a-hexahydro-3H-4,7-methanoinden-5-yl)-pyridin-3-ylmethylamine

10 g of exo-5-isothiocyanate-5,6-dihydroenedodicyclopentadiene (Maybridge international) were dissolved in 61 ml of formic acid and the solution was boiled under reflux for 45 hours. After cooling, a black precipitate was filtered off and the filtrate was concentrated. The residue was diluted with water, and 10 g of sodium hydroxide were slowly added in the presence of heat. The mixture was then cooled to room temperature, extracted three times with toluene. The combined organic phases were dried using $MgSO_4$, the $MgSO_4$ was filtered off and the filtrate was concentrated. The residue was distilled and afforded 3.38 g of a clear oil.

HPLC-RT=3.15 min; MS (CI⁺): 150 (M+H)⁺

[090] b) (exo/endo)-octahydro-4,7-methanoinden-5-ylamine

3.3 g of the double-bond isomer mixture from 1a) were dissolved in 30 ml of methanol. 0.5 g of palladium on carbon (10%), as catalyst, was added thereto and the mixture was hydrogenated under a hydrogen atmosphere for 4 h. The catalyst was then filtered off, washed with methanol and the filtrate was concentrated. After drying under a high vacuum, 3 g of product were obtained as a clear oil.

HPLC-RT=3.33 min; MS (ES⁺): 152 (M+H)⁺

LAW OFFICES


FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[091] c) exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethylamine hydrochloride

A solution of 3 g of the exo/endo-configuration octahydro-4,7-methanoinden-5-ylamine from 1 b) and 2.15 g of pyridin-3-carbaldehyde in 200 ml of toluene were heated to boiling for 5 hours in a water separator after addition of a catalytic amount of p-toluenesulfonic acid. After allowing to stand overnight at room temperature, the solvent was distilled off. The residue was dissolved in 150 ml of methanol and 0.91 g of sodium borohydride were then added in small portions to the ice-cooled solution with stirring. The mixture was stirred at room temperature for several hours and then rendered strongly acidic using excess methanolic HCl. After concentrating in a rotary evaporator, the residue was dissolved in water and rendered alkaline using potassium carbonate solution. The mixture was then extracted three times with EA, and the combined extracts were dried over magnesium sulfate, filtered, and adjusted to pH 1-2 using ethereal HCl. The solvent was then decanted off from the precipitated product and the residue was dissolved in ethanol in the warm. After cooling, the product was precipitated using ether. 2.85 g of pale crystals were obtained.

HPLC-RT=3,15, MS (ES+) : 243.2 (M+H)⁺

[092] **Example 2: (+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)-pyridin-3-ylmethylamine hydrochloride**



[093] a) (+)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine and (-)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine

The title compounds can be obtained enantiomerically pure starting from racemic (exo/endo)-octahydro-4,7-methanoinden-5-ylamine from Example 1 b) either by means

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

of chromatography on chiral columns or by crystallization using chiral acids. Using Z-valine, for example, (+)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine can be obtained as follows:

[094] a1) (+)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine by resolution using Z-valine

A mixture of 50 g of racemic (exo/endo)-octahydro-4,7-methanoinden-5-ylamine from Example 1b) and 83.2 g of Z-valine in 1.5 l of tetrahydrofuran was heated to boiling. The clear solution was allowed to cool to ambient temperature in the course of 3 hours. It was then stirred at ambient temperature for a further 20 hours. The precipitate was filtered off with suction, washed with 50 ml of tetrahydrofuran, and dried at 40°C in a drying oven. 55 g of the salt of 65% diastereomeric purity were obtained.

After recrystallizing three times from one liter of tetrahydrofuran in each case, 30 g of the salt of > 95% diastereomeric purity were obtained.

[095] 1 N sodium hydroxide solution was added to a suspension of 9.4 g of the above Z-valine salt in 30 ml of toluene and water until the pH remained constant at 11. In the course of this, the solid went into solution. The phases were separated and the water phase was extracted a further three times using 10 ml of toluene in each case. The combined toluene phases were dried over sodium sulfate, and the toluene was distilled off in vacuo.

[096] It was possible to react the amine thus obtained (3.3 g, enantiomeric purity > 95%) without further purification.

[097] a2) (-)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine by resolution using Z-D-valine:

The (-) enantiomer was obtained analogously to the above procedure using Z-D-valine. Thus, starting from 500 mg of (+)-(exo/endo)-octahydro-4,7-methanoinden-5-ylamine

from Example 1b), after precipitation and two recrystallizations, it was possible to obtain 468 mg of the salt having a diastomeric purity of > 95%.

[098] b) (+)-exo/endo-(octahydro-4,7-methanoinden-5-yl)-pyridin-3-ylmethylamine hydrochloride

A solution of 1.4 g of the (+)-exo/endo-configuration octahydro-4,7-methanoindene-5-ylamine from 2a1) and 1 g of pyridin-3-carbaldehyde in 100 ml of toluene was heated to boiling for 5 hours in a water separator after addition of a catalytic amount of p-toluenesulconic acid. After allowing the solution to stand overnight at room temperature the solvent was distilled off. The residue was dissolved in 75 ml of methanol and 0.42 g of sodium borohydride were then added in small portions to the ice-cooled solution. The mixture was stirred for several hours at room temperature, allowed to stand overnight and then rendered strongly acidic using excess methanolic HCl. The crystallization initiated by trituration was completed overnight in a refrigerator. After decanting off the solution, the residue was taken up with aqueous potassium carbonate solution and extracted three times with EA at pH 11, and the combined extracts were dried over magnesium sulfate, filtered, and adjusted to pH 1-2 using methanolic HCl. The product was then precipitated by addition of ether. 1.1 g of pale crystals were obtained.


[099] HPLC-RT=3.20; MS (CI+) : 243.3 (M+H)⁺; [α]_{Na} 589 nm: +34,6° in ethanol

2025 RELEASE UNDER E.O. 14176

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[0100] Example 3: (-)-exo/endo-(octahydro-4,7-methanoinden-5-yl)pyridin-3-ylmethanamine hydrochloride




[0101] A solution of 1.4 g of (-)-exo/endo-configuration octahydro-4,7-methanoinden-5-ylamine from 2a2) and 1 g of pyridin-3-carbaldehyde in 100 ml of toluene were heated to boiling for 5 hours in a water separator after addition of a catalytic amount of p-toluenesulfonic acid. The solvent was distilled off after allowing the solution to stand overnight at room temperature. The residue was dissolved in 75 ml of methanol and 0.42 g of sodium borohydride was then added in small portions with stirring to the ice-cooled solution. The mixture was stirred for several hours at room temperature, allowed to stand overnight and was then rendered strongly acidic using excess methanolic HCl. The crystallization initiated by trituration was completed overnight in a refrigerator. After decanting off the solution, the residue was taken up with aqueous potassium carbonate solution and extracted three times with EA at pH 11, and the combined extracts were dried over magnesium sulfate, filtered and adjusted to pH 1-2 using methanolic HCl. The product was then precipitated by addition of ether. 1.1 g of pale crystals were obtained.

[0102] HPLC-RT=3.12; MS (Cl⁺) : 243.3 (M+H)⁺; [α]_D²⁵Na 589 nm : -32.5° in ethanol

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[0103] Example 4: exo/endo-furan-2-ylmethyl-(octahydro-4,7-methanoinden-5-yl)-amine hydrochloride



[0104] 200 mg of the exo/endo-configuration octahydro-4,7-methanoinden-5-ylamines from Example 1 b), 127 mg of 2-furaldehyde, and 101 mg of p-toluenesulfonic acid were dissolved in 20 ml of toluene (anhydrous) and the solution was boiled under reflux for 4 hours. After allowing the solution to stand overnight at room temperature, the solvent was distilled off. The residue was dissolved in 15 ml of methanol and 0.6 g of sodium borohydride were then added to the ice-cooled solution in small portions with stirring. The mixture was stirred for several hours at room temperature and then rendered acidic using excess methanolic HCl. After concentrating in a rotary evaporator, the residue was taken up in 2 N NaOH and extracted three times with EA. The combined organic phases were acidified using methanolic HCl and concentrated. The oily residue was purified by means of preparative HPLC on RP-18 using acetonitrile/water (0.05% TFA). The clean fractions were combined, the acetonitrile was removed in a rotary evaporator, and the mixture was adjusted to pH 11 using potassium carbonate and extracted with EA. The combined EA phases were dried using MgSO₄ and concentrated after filtering off the MgSO₄. The residue was taken up with 2 N hydrochloric acid and freeze-dried. 119 mg of the hydrochloride were obtained as a white solid.


[0105] HPLC-RT=3.60 min; MS (ES⁺): 232.2 (M+H)⁺

FILED

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[0106] Example 5: exo/endo-(octahydro-4,7-methanoinden-5-yl)-thiophen-2-ylmethylamine hydrochloride




[0107] 200 mg of exo/endo-configuration octahydro-4,7-methanoinden-5-ylamine from Example 1b), 148 mg of thiophene-2-aldehyde and 101 mg of p-toluenesulfonic acid were dissolved in 20 ml of toluene (anhydrous) and the solution was boiled under reflux for 4 hours. After allowing the solution to stand overnight at room temperature, the solvent was distilled off. The residue was dissolved in 15 ml of methanol and 0.06 g of sodium borohydride were then added to the ice-cooled solution in small portions with stirring. The mixture was stirred for several hours at room temperature and was then rendered acidic using excess methanolic HCl. After concentrating in a rotary evaporator, the residue was taken up in 2 N NaOH and extracted three times with EA. The combined organic phases were acidified with methanolic HCl and concentrated. The oily residue was purified by means of preparative HPLC on RP-18 using acetonitrile/water (0.05% TFA). The clean fractions were combined, acetonitrile was removed in a rotary evaporator, and the mixture was adjusted to pH 11 using potassium carbonate and extracted with EA. The combined EA phases were dried using MgSO₄ and concentrated after filtering off the MgSO₄. The residue was taken up with 2 N hydrochloric acid and freeze-dried. 61 mg of the hydrochloride were obtained as a white solid.

[0108] HPLC-RT=3.84 min; MS (Cl⁺): 248.3 (M+H)⁺

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[0109] Example 6: exo/exo-(octahydro-4,7-methanoinden-5-yl)-pyridin-3-ylmethylamine hydrochloride



[0110] a) Octahydro-4,7-methano-inden-5-ole

25 g of tricyclo[5.2.1.0 (2,6)]decan-8-one (Aldrich) were dissolved in 100 ml of methanol and treated with 6.3 g of solid sodium borohydride at room temperature with slight cooling and stirring in portions in the course of 2 h. The mixture was then stirred for 2 h and allowed to stand overnight. About 40 ml of 2 N HCl were then added dropwise with cooling, followed by 20 ml of water. The mixture was concentrated, the residue was treated with ethyl acetate, and the ethyl acetate phase was washed once with water and once with sodium hydrogen carbonate solution. After drying using magnesium sulfate, the ethylacetate phase was filtered and concentrated. 26 g of oil remained, which was purified by vacuum distillation. 20.7 g of an oily liquid were obtained (bp_{0.5} 76°C).

[0111] HPLC-RT=4.55 min; MS (CI⁺): 134.8 (M-OH)⁺

[0112] b) 2-(Octahydro-4,7-methanoinden-5-yl)-isoindole-1,3-dione

1.7 g of diethyl azodicarboxylate diluted with 5 ml of THF were added with stirring to a solution of 1.66 g of octahydro-4,7-methanoinden-5-ole from 6 a), 1.47 g of phthalimide and 2.62 g of triphenylphosphine in 15 ml of THF. After standing overnight, the reaction mixture was evaporated, the residue was stirred with ether, the precipitate was filtered off with suction, and the filtrate was concentrated. The residue was purified on silica gel/toluene. 1.36 g of a yellow oil were obtained.

[0113] HPLC-RT=5.82 min; MS (CI⁺): 282.2 (M+H)⁺

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[0114] c) exo/exo-octahydro-4,7-methanoinden-5-ylamine

0.4 g of hydrazine hydrate was added dropwise to a solution of 1.12 g of 2-(octahydro-4,7-methanoinden-5-yl)-isoindole-1,3-dione from 6 b) and 15 ml of ethanol and stirred at 65°C for 2 h. The mixture was then adjusted to pH 1-2 using conc. HCl and treated with 10 ml of ethanol. The precipitate was filtered off and the filtrate was concentrated. The residue was purified by means of preparative HPLC on RP-18 using acetonitrile/water (0.05% trifluoroacetic acid). After freeze-drying, 567 mg of product were obtained as trifluoroacetate. Treatment with sodium hydroxide solution and ethyl acetate yielded 322 mg of the free amine.

[0115] HPLC-RT=3.47 min; MS (CI⁺): 152.0 (M+H)⁺

[0116] d) exo/exo-(octahydro-4,7-methanoinden-5-yl)-pyridin-3-ylmethylamine hydrochloride


A solution of 332 mg of the exo/exo-configuration octahydro-4,7-methano-indenylamine from 6 c) and 215 mg of pyridin-3-carbaldehyde in 20 ml of toluene were heated to reflux for 5 hours after addition of a catalytic amount of p-toluenesulfonic acid. After allowing the solution to stand overnight at room temperature the solvent was distilled off. The residue was dissolved in 15 ml of methanol and 91 mg of sodium borohydride were then added in small portions to the cooled solution with stirring. The mixture was stirred at room temperature for several hours and then rendered strongly acidic using excess methanolic HCl. After concentrating in a rotary evaporator, the residue was taken up with 2 N sodium hydroxide solution. After extracting three times with EA, the combined extracts were concentrated, and the residue was purified by means of preparative HPLC on RP-18 using acetonitrile/water (0.05% TFA). The product-containing fractions were combined, freeze-dried, and again purified by HPLC. The clean fractions were combined, the acetonitrile was removed on a rotary evaporator, and the residue was adjusted to pH 11 using potassium carbonate and extracted with EA. The combined EA phases were dried using MgSO₄ and concentrated after filtering off the MgSO₄. The residue was taken up with 2 N hydrochloric acid and freeze-dried. 35 mg of the hydrochloride were obtained as a white solid.

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[0117] HPLC-RT=3.25 min, MS (ES⁺): 243.1 (M+H)⁺

[0118] Example 7: exo/endo-(3-chloro-5-trifluoromethylpyridin-2-ylmethyl)(octahydro-4,7-methanoinden-5-yl)-amine trifluoroacetic acid salt




[0119] 0.5 mmol of exo/endo-octahydro-4,7-methanoindene-5-ylamine from Example 1 b), 0.5 mmol of 3-chloro-5-trifluoromethylpyridin-2-carbaldehyde, 140 μ l of triethylamine, and 10 ml of dichloromethane were introduced, 250 μ l of a 1-molar solution of titanium tetrachloride in toluene were added dropwise and the mixture was stirred at room temperature for 24 h. 1.5 ml of a 1-molar solution of sodium cyanoborohydride in THF were then slowly added and the mixture was stirred at room temperature for 30 min. The mixture was then treated with 15 ml of 2 N NaOH and stirred for 15 min. The solid was filtered off and washed with water. 30 ml of EA were added to the filtrate, the mixture was shaken and then the organic phase was separated off. After drying, the mixture was concentrated and the residue was purified by means of preparative HPLC (RP18, gradient acetonitrile/water 30% \rightarrow 90%, with 0.1% TFA in both components). After freeze-drying, 4.7 mg were obtained as a white solid.

[0120] HPLC-RT= 11.23 min*, MS (ES⁺): 345.2 (M+H)⁺

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[0121] Example 8: *exo/endo*-(decahydro-1,4-methanonaphthalen-2-yl)-pyridin-3-ylmethyl-amine hydrochloride



[0122] a) bis-(3-chloro-1,2,3,4-tetrahydro-1,4-methanonaphthalen-2-yl)-diazene N,N'-dioxide

3.34 g of isoamyl nitrite were added to a solution of 3.56 g of benzonorbornadiene [L. Friedman and F.M. Logullo, J. Org. Chem. 34: 3089-3092, (1969)] in 6 ml of glacial acetic acid and 6 ml of ethanol. 8.5 ml of a 15% strength solution of hydrogen chloride gas in ethanol were then added dropwise. The resulting suspension was stirred at room temperature for a further 2½ hours and then treated with 20 mg of diisopropyl ether. The solid was filtered off after further stirring for 30 minutes. A pale crystalline solid; m.p. 187 - 188°C, was obtained.

MS (FAB): 415.1 (M+H)⁺

[0123] b) (*exo*)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-2-ylamine

3 g of bis-(3-chloro-1,2,3,4-tetrahydro-1,4-methanonaphthalen-2-yl)-diazene N,N'-dioxide were suspended in 150 ml of methanol and hydrogenated in an autoclave with hydrogen at 100 bar and 100°C for 20 hours using Raney nickel catalyst. After filtering off the catalyst, the solvent was evaporated. The residue was treated with water, rendered strongly alkaline with NaOH, and extracted repeatedly with methyl tert-butyl ether. After drying the organic phases, the desired amine was obtained as a pale yellow liquid.

[0124] MS (ES⁺): 160.0 (M+H)⁺

[0125] c) exo/endo-decahydro-1,4-methanonaphthalen-2-ylamine

A solution of 1 g of exo/endo-1,2,3,4-tetrahydro-1,4-methanonaphthalen-2-ylamine in 10 ml of methanol and 30 ml of 2 N hydrochloric acid was hydrogenated in an autoclave with hydrogen at 100 bar and 90°C for 10 hours using 0.4 g of RuO₂. After separating off the catalyst, the solution was evaporated to half the original volume. The aqueous solution thus obtained was rendered strongly alkaline with 10 N NaOH and extracted repeatedly with methyl tert-butyl ether. After drying and evaporating the solvents, exo/endo-decahydro-1,4-methanonaphthalen-2-ylamine was obtained as a colorless oil, which was stored under argon.

[0126] MS (CI⁺): 166.2 (M+H)⁺

[0127] d) exo/endo-(decahydro-1,4-methanonaphthalen-2-yl)-pyridin-3-ylmethylamine hydrochloride

A solution of 0.71 g of pyridin-3-aldehyde and 1.1 g of exo/endo-decahydro-1,4-methanonaphthalene-2-ylamine in 40 ml of toluene was boiled under reflux for 4 hours after addition of a small catalytic amount of p-toluenesulfonic acid (3-5 mg) and the solvent was then distilled off. After dissolving the oily residue in about 30 ml of anhydrous methanol, the solution was treated in portions with 0.335g of sodium borohydride with cooling and stirring, stirred at room temperature for a further 2 hours, and allowed to stand overnight. The solution was then rendered acidic using a solution of HCl in methanol, the precipitate was filtered off, and the solvent was evaporated. The residue was recrystallized from ethanol, m.p. 283-285°C.







[0128] HPLC-RT=3.38 min, MS (CI⁺): 257.4 (M+H)⁺

[0129] The compounds described below were prepared from carbonyl derivatives known from the literature and the appropriate amines, analogously to the example indicated:

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000








Table 1

Example		Salt	Analogously to Example	MS	HPLC RT [min]
9		HCl	5	ES+ 232.2 (M+H)+	3.73
10		HCl	5	ES+ 260.2 (M+H)+	4.00
11		HCl	5	Cl+ 248.0 (M+H)+	3.88
12		HCl	5	ES+ 243.1 (M+H)+	3.63
13		HCl	5	Cl+ 243.0 (M+H)+	3.14
14		HCl	5	Cl+ 232.2 (M+H)+	3.05








202-408-4000

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

15		HCl	5	Cl+ 232.1 (M+H)+	3.05
16		HCl	5	ES+ 231.2 (M+H)+	3.76
17		HCl	5	ES+ 249.1 (M+H)+	3.47
18 (racemate)		HCl	5	Cl+ 244.1 (M+H)+	3.39
19 (+)		HCl	2	Cl+ 244.2 (M+H)+	3.41
20 (-)		HCl	3	Cl+ 244.2 (M+H)+	3.40
21		HCl	8	ES+ 258.1 (M+H)+	3.65




100-408-4000

22		HCl	5	ES+ 244.1 (M+H)+	3.33
23		HCl	5	ES+ 244.2 (M+H)+	3.43
24		HCl	5	Cl+ 232.1 (M+H)+	3.42
25	 mixture 	HCl	5	Cl+ 241.2 (M+H)+	3.07
26		TFA	7	ES+ 330.1 (M+H)+	11.78*
27		TFA	7	ES+ 274.2 (M+H)+	9.75*

202-408-4000

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

28		TFA	7	ES+ 293.1 (M+H)+	10.03*
29		TFA	7	ES+ 356.3 (M+H)+	11.18*
30		TFA	7	ES+ 319.3 (M+H)+	9.99*

PHARMACOLOGICAL DATA

Example 31: Description of the Caco 2 model

[0130] The Caco-2 cell line was acquired from the American Type Culture Collection (ATCC) and maintained in Dulbecco's Modified Eagle medium (high glucose content), supplemented with nonessential amino acids, L-glutamine, penicillin/streptomycin, and 10% strength fetal calf serum. The Caco-2 cell line was then kept in an incubator under a 10% strength CO₂ atmosphere at 95% strength relative humidity and 37°C. The cells were grown in cell culture flasks (175 cm²). For the transport investigations, the Caco-2 cells were inoculated into polycarbonate cell culture inserts (costar Transwells®, pore size: 3 µm, area: 4.71 cm²) at a cell density of 6.5 x 10⁴ cells/cm² and incubated in six-well culture plates with a change of medium after 4 and 8 days and then every second day thereafter. 21 to 25 day-old monolayers were used for the experiments.

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[0131] In each test series, a 21 day-old monolayer was tested for its properties with ³H-dextran as a permeability marker. The value of the transfer rate (cumulative) had to be in the range of 2% after 120 min.

[0132] After eliminating the growth medium from the apical and the basolateral side, the monolayers were rinsed with the transport buffer (Hank's balanced salt solution, pH 7.8; containing 2.8 g/l glucose). The cells were then equilibrated for 15 min at 37°C under a 10% strength CO₂ atmosphere. The HBSS buffer was then removed. The test compounds were dissolved in a mixture of HBSS buffer and DMSO and added to the apical buffer such that a 1% strength (v/v) DMSO solution resulted. The test concentration in the first experiment was 1 mM, and 100 µM in the second. The experiments were carried out at 37°C and started by the addition of 1.5 ml of test solution to the donor side (apical). Transport buffer without compound was added to the recipient side (basolateral, 2.5 ml). At various points in time, samples were taken from the basolateral side (1 ml) and replaced by fresh buffer solution at 37°C. Apical samples were taken at the start and at the end (120 min) in order to determine the recovery rate of the compounds by means of these concentrations and the cumulative basolateral concentration.

[0133] The compounds were analyzed by means of HPLC.

[0134] The apparent permeability coefficient (P_{app}) was calculated by means of the following equation:

$$P_{app} = \frac{d_c \cdot V}{d_t \cdot A \cdot c_o}$$

[0135] wherein d_c/d_t is the flow through the monolayer (µg of compound/ml x s), V is the liquid volume in the collection chamber (ml), A is the surface area size of the monolayer (cm²) and c_o is the initial concentration (µg of compound/ml) in the donor chamber. The flow through the monolayer was calculated from the cumulative basolateral concentration at the appropriate point in time with the aid of the initially linear data curve (linear up to 60 min). The respective determinations were made in





LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[illegible]

Results:

[0136] Table 2

		Absorbed fraction [%]
Example 1		100
Example 18		100
S 3226		<5
S 2120		<1

2025 RELEASE UNDER E.O. 14176

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

[0137] Compared with the NHE3 active compounds of the acylguanidine type known from the literature (J.-R. Schwark et al. Eur. J. Physiol., 436:797 (1998)), the compounds of the formula (I) or (I a) show a clearly superior ability to cross the membrane.

Example 32: Description of the NHE activity measurements

[0138] Most of the molecular biology techniques follow protocols from the works "Current Protocols in Molecular Biology" (Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A. and Struhl, K., eds., John Wiley & Sons) and "Molecular Cloning: A Laboratory Manual" (Sambrock, J., Fritsch, E.F., and Maniatis, T., Cold Spring Harbor Laboratory Press (1989)).

[0139] In the context of our studies, stably transfected cell lines were produced that in each case expressed one of the following NHE subtypes: Human NHE1 (Sardet *et al.* Cell 56:271-280 (1989)), rabbit NHE2 (Tse *et al.* J. Biol. Chem. 268:11917-11924(1993)), human NHE3 (Brant *et al.* Am. J. Physiol. 269 (Cell Physiol. 38) C198-C206 (1995)), or rat NHE3 (Orlowski *et al.*; J. Biol. Chem. 267:9331-9339 (1992)).

[0140] The cDNA clones of the respective NHE subtypes obtained by Prof. Pouyssegur were, after addition of suitable linker sequences, cloned into the expression plasmid pMAMneo (obtainable, for example, via CLONTECH, Heidelberg) such that the recognition sequence for the restriction endonuclease NheI of the plasmid was approximately 20 – 100 Basepairs before the start codon of the respective NHE subtype and that the total coding sequence was present in the construct. In the case of the human NHE3 obtained from human kidney mRNA by means of RT-PCR, the RT-PCR primers were selected such that the cDNA band obtained had cleavage sites suitable for pMAMneo at its ends.

[0141] Using the "calcium phosphate method" (described in chapter 9.1 of "Current Protocols in Molecular Biology"), the NHE-deficient cell line LAP1 (Franchi *et*

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

al.; Proc. Natl. Acad. Sci. USA 83:9388-9392 (1986)) was transfected with the plasmids that received the respective coding sequences of the NHE subtypes. After selection for transfected cells by means of growth in G418-containing medium (only cells which have obtained a neogene by transfection can survive under these conditions), selection was made for functional NHE expression. For this, the "Acid Load" technique described by Sardet was used (Sardet *et al.*; Cell 56:271-280 (1989)). Unlike untransfected LAP1 cells, cells that express a functional NHE subtype can also compensate for the acidification carried out in this test in the absence of CO_2 and HCO_3^- . After repetition of the "Acid Load" selection several times, the surviving cells were inoculated into microtiter plates such that one cell per well should occur statistically. Under the microscope, after approximately 10 days, a check was made to estimate how many colonies were growing per well. Cell populations from individual colonies were then investigated using the XTT proliferation kit (Boehringer Mannheim) with respect to their survival ability after "Acid Load". The best cell lines were used for further tests and, to avoid a loss of the transfected sequence, cultured in G418-containing medium under continuous selection pressure.

[0142] To determine IC_{50} values for the inhibition of the individual NHE subtypes by specific substances, a test developed by S. Faber (Faber *et al.*; Cell. Physiol. Biochem. 6:39-49 (1996)), which is based on the "Acid Load" technique, was slightly modified.

[0143] In this test, the recovery of the intracellular pH (pH_i) after an acidification, which in the case of functional NHE occurs even under bicarbonate-free conditions, was determined. For this, the pH_i was determined using the pH-sensitive fluorescent dye BCECF (Calbiochem, where the precursor BCECF-AM is employed). The cells were first loaded with BCECF. The BCECF fluorescence was determined in a "Ratio Fluorescence Spectrometer" (Photon Technology International, South Brunswick, N.J., USA) at excitation wavelengths of 505 and 440 nm and an emission wavelength of

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

535 nm and converted into the pH_i by means of calibration curves. Differing from the described protocol, the cells were incubated in NH_4Cl buffer (pH 7.4) even in the case of the BCECF loading (NH_4Cl buffer: 115 mM NaCl, 20 mM NH_4Cl , 5 mM KCl, 1 mM CaCl_2 , 1 mM MgSO_4 , 20 mM Hepes, 5 mM glucose, 1 mg/ml BSA; a pH of 7.4 is established using 1 M NaOH). The intracellular acidification was induced by addition of 975 μl of an NH_4Cl -free buffer to 25 μl aliquots of the cells incubated in NH_4Cl buffer. The rate of pH recovery that followed acidification was recorded for 2 minutes in the case of NHE1, for 5 minutes in the case of NHE2, and for 3 minutes in the case of NHE3. For the calculation of the inhibitory potency of the tested substances, cells were first investigated in buffers in which a complete pH recovery occurred and in buffers in which no pH recovery at all took place. For the complete pH recovery (100%), the cells were incubated in Na^+ -containing buffer (133.8 mM NaCl, 4.7 mM KCl, 1.25 mM CaCl_2 , 1.25 mM MgCl_2 , 0.97 mM Na_2HPO_4 , 0.23 mM NaH_2PO_4 , 5 mM hepes, 5 mM glucose, a pH of 7.0 is established using 1 M NaOH). For the determination of the 0% value, the cells were incubated in an Na^+ -free buffer (133.8 mM choline chloride, 4.7 mM KCl, 1.25 mM CaCl_2 , 1.25 mM MgCl_2 , 0.97 mM K_2HPO_4 , 0.23 mM KH_2PO_4 , 5 mM hepes, 5 mM glucose, a pH of 7.0 is established using 1 M NaOH). The substances to be tested were prepared in the Na^+ -containing buffer. The recovery of the intracellular pH at any tested concentration of a substance was expressed as a percent of the maximum recovery. From the percentage values of the pH recovery, the IC_{50} value of the respective substance was calculated by means of a Sigma Plot program for the individual NHE subtypes.

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000

NHE activity

Table 3

Example	Rat NHE3 IC ₅₀ [μM]
1	0.34
5	2.9
6	2.1

11/11/2011 11:11:11 AM

LAW OFFICES

FINNEGAN, HENDERSON,
FARABOW, GARRETT,
& DUNNER, L.L.P.
1300 I STREET, N. W.
WASHINGTON, DC 20005
202-408-4000